

Growth and photosynthesis of *Chlorella* strains from polar, temperate and tropical freshwater environments under temperature stress*

Kok-Keong LEE^{1,2}, Phaik-Eem LIM^{1,**}, Sze-Wan POONG¹, Chiew-Yen WONG^{3,4}, Siew-Moi PHANG^{1,5}, John BEARDALL⁶

¹ Institute of Ocean and Earth Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

² Institute of Graduate Studies, University of Malaya, 50603 Kuala Lumpur, Malaysia

³ School of Health Sciences, International Medical University, 57000 Kuala Lumpur, Malaysia

⁴ National Antarctic Research Centre, University of Malaya, 50603 Kuala Lumpur, Malaysia

⁵ Institute of Biological Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

⁶ School of Biological Sciences, Monash University, Clayton, Victoria, 3800, Australia

Received Mar. 28, 2017; accepted in principle May 8, 2017; accepted for publication Jun. 21, 2017

© Chinese Society for Oceanology and Limnology, Science Press, and Springer-Verlag GmbH Germany 2017

Abstract Elevated temperatures as a consequence of global warming have significant impacts on the adaptation and survival of microalgae which are important primary producers in many ecosystems. The impact of temperature on the photosynthesis of microalgae is of great interest as the primary production of algal biomass is strongly dependent on the photosynthetic rates in a dynamic environment. Here, we examine the effects of elevated temperature on *Chlorella* strains originating from different latitudes, namely Antarctic, Arctic, temperate and tropical regions. Chlorophyll fluorescence was used to assess the photosynthetic responses of the microalgae. Rapid light curves (RLCs) and maximum quantum yield (F_v/F_m) were recorded. The results showed that *Chlorella* originating from different latitudes portrayed different growth trends and photosynthetic performance. The *Chlorella* genus is eurythermal, with a broad temperature tolerance range, but with strain-specific characteristics. However, there was a large overlap between the tolerance range of the four strains due to their “eurythermal adaptivity”. Changes in the photosynthetic parameters indicated temperature stress. The ability of the four strains to reactivate photosynthesis after inhibition of photosynthesis under high temperatures was also studied. The *Chlorella* strains were shown to recover in terms of photosynthesis and growth (measured as Chl *a*) when they were returned to their ambient temperatures. Polar strains showed faster recovery in their optimal temperature compared to that under the ambient temperature from which they were isolated.

Keyword: Antarctic; Arctic; F_v/F_m ; microalgae; pigments; recovery

1 INTRODUCTION

Anthropogenic influences have led to significant changes in climate. The average global temperature has risen by about 0.8°C since 1880 and the average global surface temperature is expected to increase by 4–5°C over the next century (IPCC, 2007). In addition to higher average temperatures, global climate change is also resulting in higher temperature variability, thus increasing the risks to species’ tolerance limits. Temperature is one of the major factors affecting the distribution and productivity of microalgae.

Understanding the growth and photosynthetic response of microalgae to the changes in their thermal environment is crucial to the characterization of their ecophysiology in nature. Temperature effects on the

* Supported by the HiCoE Grant (No. IOES-2014H) from the Ministry of Higher Education (MOHE), Malaysia, the University of Malaya Postgraduate Research Fund (No. PG121-2015A), the Knowledge Management Grant (No. RP001O-13SUS), and the Antarctic Flagship Project (Nos. FP0712E012, PV002-2015) by the Ministry of Science, Technology & Innovation (MOSTI), Malaysia

** Corresponding author: phaikem@um.edu.my