



Purification and identification of novel cytotoxic oligopeptides from soft coral *Sarcophyton glaucum*^{*#}

Yixian QUAH¹, Nor Ismaliza MOHD ISMAIL^{2,3}, Jillian Lean Sim OOI⁴, Yang Amri AFFENDI^{5,6},
Fazilah ABD MANAN⁷, Lai-Kuan TEH^{3,8}, Fai-Chu WONG^{1,3}, Tsun-Thai CHAI^{†‡1,3}

¹Department of Chemical Science, Faculty of Science, Universiti Tunku Abdul Rahman, 31900 Kampar, Malaysia

²Department of Biological Science, Faculty of Science, Universiti Tunku Abdul Rahman, 31900 Kampar, Malaysia

³Centre for Biodiversity Research, Universiti Tunku Abdul Rahman, 31900 Kampar, Malaysia

⁴Department of Geography, Faculty of Arts and Social Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

⁵Institute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

⁶Institute of Ocean and Earth Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

⁷Department of Biosciences, Faculty of Science, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Malaysia

⁸Department of Biomedical Science, Faculty of Science, Universiti Tunku Abdul Rahman, 31900 Kampar, Malaysia

[†]E-mail: chaitt@utar.edu.my

Received Dec. 1, 2017; Revision accepted Apr. 19, 2018; Crosschecked Dec. 5, 2018


Abstract: Globally, peptide-based anticancer therapies have drawn much attention. Marine organisms are a reservoir of anticancer peptides that await discovery. In this study, we aimed to identify cytotoxic oligopeptides from *Sarcophyton glaucum*. Peptides were purified from among the *S. glaucum* hydrolysates produced by alcalase, chymotrypsin, papain, and trypsin, guided by a methylthiazolyldiphenyl-tetrazolium bromide (MTT) assay on the human cervical cancer (HeLa) cell line for cytotoxicity evaluation. Purification techniques adopted were membrane ultrafiltration, gel filtration chromatography, solid phase extraction (SPE), and reversed-phase high-performance liquid chromatography (RP-HPLC). Purified peptides were identified by de novo peptide sequencing. From papain hydrolysate, three peptide sequences were identified: AGAPGG, AERQ, and RDTQ (428.45, 502.53, and 518.53 Da, respectively). Peptides synthesized from these sequences exhibited cytotoxicity on HeLa cells with median effect concentration (EC₅₀) values of 8.6, 4.9, and 5.6 mmol/L, respectively, up to 5.8-fold stronger than the anticancer drug 5-fluorouracil. When tested at their respective EC₅₀, AGAPGG, AERQ, and RDTQ showed only 16%, 25%, and 11% cytotoxicity to non-cancerous Hek293 cells, respectively. In conclusion, AERQ, AGAPGG, and RDTQ are promising candidates for future development as peptide-based anticancer drugs.

Key words: Anticancer therapy; Bioactive peptide; Cytotoxicity; HeLa cells; *Sarcophyton glaucum*; Soft coral
<https://doi.org/10.1631/jzus.B1700586> **CLC number:** Q516

[‡] Corresponding author

^{*} Project supported by the Fundamental Research Grant Scheme of the Ministry of Higher Education, Malaysia (FRGS/1/2013/ST04/UTAR/02/1)

[#] Electronic supplementary materials: The online version of this article (<https://doi.org/10.1631/jzus.B1700586>) contains supplementary materials, which are available to authorized users

 ORCID: Tsun-Thai CHAI, <https://orcid.org/0000-0003-3716-1599>

© Zhejiang University and Springer-Verlag GmbH Germany, part of Springer Nature 2019

1 Introduction

Cancer is one of the leading causes of morbidity and mortality worldwide. There were about 14 million new cancer cases and 8.2 million cancer deaths in 2012 globally. It was projected that in the next 20 years, new cases will increase by about 63% (Ferlay et al., 2018). Unfortunately, chemotherapy, a common cancer treatment, often incurs side effects, including nausea,